

The following SIG codes are standard in PharmaClik Rx. Additional codes can be customized in List Maintenance (visit [PharmaClik Rx Online Help](#) for instructions).

PC	After Meals	AU	In Each Ear
APL	Applicatorful	OU	In Each Eye
APP	Apply	IEN	In Each Nostril
AR	Arthritis	AL	In the Left Ear
UD	As Directed	AD	In the Right Ear
HS	At Bedtime	OS	In the Left Eye
STAT	At Once	OD	In the Right Eye
		INF	Inflammation
BP	Blood Pressure	I	Inhale
AC	Before Meals	INJ	Inject
BM	Bowel Movement	MEDAP	Injection Administered by Pharmacist
PO	By Mouth	IR	Insert
		INS	Instil
CAP	Capsule	R	Into the Rectum
CAPS	Capsules	V	Into the Vagina
CI	Circulation		
CON	Constipation	JU	Juice
MEDCP	Consultation Provided By		
CR	Crush	N	Night
DY	Daily	SS	One Half
D	Day	QBID	One or Two Times Daily
DS	Days		
DI	Diarrhea	PA	Pain
DR	Drink	PL	Place
GTT	Drop	PF	Puffs
GTTS	Drops		
		RA	Rash
QD	Each Day	MEDRA	Refer to Adaptation Form
QPM	Each Evening	MEDRO	Refer to Opinions Form
QAM	Each Morning		
Q	Every	SW	Shake Well
Q1H	Every Hour	SB	Shortness of Breath
Q2D	Every Second Day	SP	Sparingly
Q2H	Every 2 Hours	SUPP	Suppository
Q3H	Every 3 Hours		
Q4H	Every 4 Hours	TBL	Tablespoon
Q6H	Every 6 Hours	TBLS	Tablespoonfuls
Q8H	Every 8 Hours	TAB	Tablet
Q12H	Every 12 Hours	TABS	Tablets
Q2-3H	Every 2-3 Hours	T	Take
Q2-4H	Every 2-4 Hours	TSP	Teaspoon
Q4-6H	Every 4-6 Hours	TSPS	Teaspoonfuls
		TQID	Three or Four Times Daily
FE	Fever	TID	Three Times Daily
FL	Fluids	BTID	Two or Three Times Daily
F	For	ADMX	To a Maximum of
QID	Four Times Daily	AA	To Affected Area
		BID	Twice Daily
G	Give		
		SL	Under the Tongue
HD	Headache	UF	Until Finished
HR	Heart		
H	Hour	AQ	Water
HRS	Hours	PRN	When Required
		PRNF	When Necessary For
		C	With
		CF	With Food
		CC	With Meals
		S	Without